

Abstract Submitted  
for the MAR10 Meeting of  
The American Physical Society

**Effects of Nanoparticle Size Polydispersity On the Tethered Nanoparticle Phase Diagram** CAROLYN PHILLIPS, SHARON GLOTZER, University of Michigan — Recent simulations predict that aggregating nanospheres functionalized with polymer “tethers” can self-assemble to form a cylinder, perforated lamellae, lamellae, and even the double gyroid phase also seen in block copolymer and surfactant systems [1]. We study the impact of nanoparticle size polydispersity on the properties of the phase diagram[2],[3]. We show that in the portions of the phase diagram characterized by an icosahedral packing motif, a low amount of polydispersity lowers the energy and a large amount of polydispersity raises the potential energy of the system by disrupting the icosahedral packing. In the portions of the phase diagram characterized by crystalline packing, polydispersity raises the energy of the system and induces a phase transition from crystalline to liquid within the nanosphere packing of the microphase.

[1] Iacovella, et al., PRE, 2007

[2] Phillips, et al., “Stability of the double gyroid phase to nanoparticle polydispersity in polymer tethered nanosphere systems, preprint.

[3] Phillips, et al., “Effects of Nanoparticle Size Polydispersity On the Tethered Nanoparticle Phase Diagram,” preprint.

Carolyn Phillips  
University of Michigan

Date submitted: 28 Nov 2009

Electronic form version 1.4