

Abstract Submitted
for the MAR10 Meeting of
The American Physical Society

Transient XUV lasing without inversion via He triplet states¹

E.A. SETE, Department of Physics and Institute for Quantum Studies, Texas A&M University, S. SUCKEWER, Department of Mechanical & Aerospace Engineering, Princeton University, A.A. SVIDZINSKY, Department of Physics and Institute for Quantum Studies, Texas A&M University, M.O. SCULLY, Department of Physics and Institute for Quantum Studies, Texas A&M University; Princeton University — We demonstrate transient inversionless laser oscillation in short wavelength regions where population inversion is difficult to achieve. It was shown some time ago that when the excitation of an ensemble of two-level atoms is swept in the direction of lasing, so that atoms are prepared in the excited state just as the radiation from previously excited atoms reaches them, the resulting laser amplifier is “highly anomalous” and yields superradiant emission without population inversion. We here show that transient lasing without inversion in a three-level system has common features with Dicke superradiance and can yield strong XUV lasing in, for example, He atoms (at 58 nm) or He-like ions such as B³⁺ (at 6.1 nm).

¹We gratefully acknowledge support of the Office of Naval Research (N00014-07-1-1084, N00014-08-1-0948, and N00014-09-1-0888), the Robert A. Welch Foundation (Award A-1261), and NSF Grant EEC-0540832 (MIRTHE ERC).

Eyob Sete
Dept of Physics and Institute for Quantum Studies,
Texas A&M University

Date submitted: 02 Dec 2009

Electronic form version 1.4