

Abstract Submitted  
for the MAR11 Meeting of  
The American Physical Society

**Anisotropy Energy, Spin-Atomic Vibration Interaction, and Spin-Flip Hamiltonian of a Single Atomic Spin in a Crystal Field** SATOSHI KOKADO, Faculty of Engineering, Shizuoka University, Japan, KIKUO HARI-GAYA, Nanosystem Research Institute, AIST, Japan, AKIMASA SAKUMA, Graduate School of Engineering, Tohoku University, Japan — We derive the anisotropy energy  $V_A$ , the spin-atomic vibration interaction  $V_{SA}$ , and the spin-flip Hamiltonian  $V_{SF}$  of a single atomic spin system, “ $\text{Fe}^{2+}$  ( $3d^6$ ) in a crystal field of tetragonal symmetry” [1,2]. We here apply the perturbation theory to a model with the spin-orbit interaction and the kinetic and potential energies of electrons. The model also takes into account the difference in vibration displacement between an effective nucleus and electrons,  $\Delta r$ . We first find conditions to enhance a coefficient  $|D|$  of  $V_A = -|D|S_Z^2$ , where  $D$  is an anisotropy constant and  $S_Z$  is the  $Z$  component of a spin operator. Second, we show that  $V_{SA}$  appears for  $\Delta r \neq 0$ , while  $V_{SF}$  is present independently of  $\Delta r$ . Also, the magnitudes of the coefficients of  $V_{SA}$  can be larger than those of the conventional spin-phonon interaction depending on vibration frequency.

[1] S. Kokado *et al.*, J. Phys. Soc. Jpn. **79**, 114721 (2010).

[2] S. Kokado *et al.*, phys. stat. solidi (c) **7**, 2612 (2010).

Satoshi Kokado  
Faculty of Engineering, Shizuoka University

Date submitted: 15 Nov 2010

Electronic form version 1.4