

Abstract Submitted  
for the MAR11 Meeting of  
The American Physical Society

**Micellization and phase transitions in a triblock copolymer-D<sub>2</sub>O system**<sup>1</sup> HOSANNA ODHNER, Bryn Mawr, ALISON HUFF, KELLY PATTON, D.T. JACOBS, The College of Wooster, BRYNA CLOVER, University of Maryland, College Park, SANDRA GREER<sup>2</sup>, University of Maryland and Mills College — The triblock copolymer (“unimer”) of PPO-PEO-PPO (commercially known as 17R4) has hydrophobic ends and a hydrophilic center. When placed in D<sub>2</sub>O at lower concentrations and temperatures, only a network of unimers exists. However, at higher concentrations or temperatures, micelles of different geometries can form. We have measured the micellization line marking the transition from only unimers to some micelles, as well as a one- to two-phase transition at higher temperatures. This second transition is an Ising-like, LCST critical point, based on the shape of the coexistence curve. We find the LCST to not correspond to the minimum of the cloud point curve, which indicates polydispersity as described by Sollich.

<sup>1</sup>We acknowledge the support from Research Corporation, NSF-REU grant DMR 0649112, The College of Wooster, and (for BC and SG) to the donors of the Petroleum Research Fund, administered by the American Chemical Society

<sup>2</sup>currently Provost at Mills College

Donald Jacobs  
The College of Wooster

Date submitted: 17 Nov 2010

Electronic form version 1.4