

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

Variational Study of a Finite Heisenberg Chain ERIC ASHENDORF, JOE WEINER, JAY D. MANCINI, Kingsborough College of CUNY, VASSILIOS FESSATIDIS, Fordham University, SAMUEL P. BOWEN, Chicago State University — Here we wish to study the ground-state of the 1D Heisenberg chain

$$H = -\frac{1}{2}J \sum_{l=1}^N \left[2 \left(\sigma_l^+ \sigma_{l+1}^+ + \sigma_l^- \sigma_{l+1}^- \right) + \sigma_l^z \sigma_{l+1}^z \right],$$

where the σ 's are the usual Pauli spin matrices and J is the strength of the spin-spin interaction. The purpose of our revisiting such a well known system is to use it as a benchmark for our variational ansatz in which a trial vector is chosen $|\psi_0(\alpha)\rangle = \exp\left(\alpha \sum_{l=1}^N \sigma_l^+ \sigma_{l+1}^z\right) |0\rangle_N$, where α is the variational parameter and $|0\rangle_N$ is an appropriately chosen initial array of spins. We then construct a basis according to the prescription $|\psi_m\rangle = \partial_\alpha^m |\psi_0(\alpha)\rangle$ creating an energy matrix with elements $h_{ij} = h_{ij}(\alpha, J)$ whose eigenvalues are then evaluated.

Vassilios Fessatidis
Fordham University

Date submitted: 16 Nov 2010

Electronic form version 1.4