

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

First-principles study of the noncentrosymmetric superconductors $\text{Li}_2\text{Pt}_3\text{B}$ and $\text{Li}_2\text{Pd}_3\text{B}$ TATSUYA SHISHIDOU, ADSM, Hiroshima Univ, TAMIO OGUCHI, ISIR, Osaka Univ — Effect of spin-orbit coupling (SOC) associated with lack of space-inversion symmetry has been a central issue in condensed-matter physics. $\text{Li}_2\text{Pt}_3\text{B}$ and $\text{Li}_2\text{Pd}_3\text{B}$ are superconducting below 2.7 K and 7 K, respectively, and have the same crystal structure (cubic $P4_332$, No. 212), which is non-centrosymmetric and is characterized by highly distorted corner-sharing BPt(Pd)_6 octahedra. Despite the same valency and structure, they show quite different superconducting behavior. NMR measurements [1] indicate that $\text{Li}_2\text{Pt}_3\text{B}$ would be a spin-triplet superconductor with line nodes in the gap function while $\text{Li}_2\text{Pd}_3\text{B}$ is a conventional spin-singlet s -wave superconductor. SOC would be a key to understand this difference. To clarify the electronic band structure and Fermi surface of these compounds, we performed density-functional (GGA PBE) calculations with FLAPW method. Relativistic effects were fully taken into account. The band structures calculated are in good accordance with previous work[2] and the spin splitting due to SOC is quite significant in $\text{Li}_2\text{Pt}_3\text{B}$. This work was supported by a MEXT KAKENHI on Innovative Areas “Topological Quantum Phenomen”. [1] M. Nishiyama, Y. Inada, and G.-q. Zheng, PRL **98**, 047002 (2007); PRB **71**, 220505(R) (2005). [2] K.-W. Lee and W. E. Pickett, PRB **72**, 174505 (2005).

Tatsuya Shishidou
ADSM, Hiroshima University

Date submitted: 17 Nov 2010

Electronic form version 1.4