

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

Measurement of the statistical properties of the persistent current in normal metal rings MANUEL CASTELLANOS BELTRAN, WILL SHANKS, DUSTIN NGO, Yale University, ANIA BLESZYNSKI-JAYICH, UCSB, JACK HARRIS, Yale University — A striking manifestation of quantum mechanics at the mesoscopic scale is the existence of an equilibrium persistent current in normal metal rings threaded by a magnetic flux. A theory of non-interacting diffusive electrons predicts that the amplitude of these currents is a stochastic function of the disorder profile of the specific ring. Thus the persistent current is different from sample to sample, with a Gaussian distribution. Due to the difficulty of measuring these currents, experiments to determine the form of the persistent current distribution had not yet been performed. However, our group recently developed a technique for measuring persistent currents in normal metal rings with high SNR, low measurement back-action, excellent background rejection, and over a large range of magnetic fields. We have measured a total of roughly 100 independent realizations of persistent current amplitudes in single rings. Within the statistical limits of our data, we corroborate that the first five cumulants are consistent with a Gaussian distribution. As a further test of the higher-order statistical properties of the persistent current, we also show that the quadrature amplitudes of the current's Aharonov-Bohm oscillations are uncorrelated.

Manuel Castellanos Beltran
Yale University

Date submitted: 14 Dec 2010

Electronic form version 1.4