

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

Origin of colossal ionic conductivity in YSZ/STO multilayers

TIMOTHY PENNYCOOK, MARK OXLEY, Vanderbilt Univ., Oak Ridge National Laboratory, MATTHEW BECK, Vanderbilt Univ., JAVIER GARCIA-BARRIOCANAL, FLAVIO BRUNO, CARLOS LEON, JACOBO SANTAMARIA, Complutense Univ., Madrid, MARIA VARELA, Oak Ridge National Laboratory, STEPHEN PENNYCOOK, Oak Ridge National Laboratory, Vanderbilt Univ., SOKRATES PANTELIDES, Vanderbilt Univ., Oak Ridge National Laboratory — A colossal eight order of magnitude increase in the ionic conductivity of yttria-stabilized zirconia (YSZ) near room temperature was recently reported in YSZ/strontium titanate (STO) epitaxial heterostructures [1]. We present density functional theory results that explain the enhancement in terms of strain- and interface-induced disorder of the YSZ O-sublattice [2]. We further present experimental confirmation of O disorder using a combination of scanning transmission electron microscopy and electron energy loss spectroscopy. The O K-edge fine structure shows blurred-out features indicative of disorder [3]. Atomic-resolution elemental mapping clearly resolves the O sublattice in the STO but is blurred out in the YSZ, indicating disorder. This work is supported by DOE grant DE-FG02-09ER46554 and DOE Materials Sciences and Engineering Div. 1 J. Garcia-Barriocanal et al. *Science* 321, 676 (2008); 2 T.J. Pennycook et al., *Phys. Rev. Lett.* 104, 115901 (2010); 3 T.J. Pennycook et al., *European Phys. J.*

Timothy Pennycook
Vanderbilt Univ., Oak Ridge National Laboratory

Date submitted: 17 Nov 2010

Electronic form version 1.4