

Abstract Submitted  
for the MAR11 Meeting of  
The American Physical Society

**On the critical behaviour of two-dimensional liquid crystals ANA**

I. FARIÑAS-SÁNCHEZ, Departamento de Física, Universidad Simón Bolívar, Venezuela, ROBERT BOTET, Laboratoire de Physique des Solides, Université Paris-Sud, France, BERTRAND BERCHE, Groupe de Physique Statistique, Institut Jean Lamour, France, RICARDO PAREDES, Laboratorio de Simulación, Instituto de Matemáticas. Unidad Cuernavaca. UNAM, Mexico — The Lebwohl-Lasher (LL) model is the traditional model used to describe the nematic-isotropic transition of real liquid crystals. In this paper, we develop a numerical study of the temperature behaviour and of finite-size scaling of the two-dimensional (2D) LL-model. We discuss two possible scenarios. In the first one, the 2D LL-model presents a phase transition similar to the topological transition appearing in the 2D XY-model. In the second one, the 2D LL-model does not exhibit any critical transition, but its low temperature behaviour is rather characterized by a crossover from a disordered phase to an ordered phase at zero temperature. We realize and discuss various comparisons with the 2D XY-model and the 2D Heisenberg model. Having added finite-size scaling behaviour of the order parameter and conformal mapping of order parameter profile to previous studies, we analyze the critical scaling of the probability distribution function; hyperscaling relations and stiffness order parameter and conclude that the second scenario (no critical transition) is the most plausible.

Ana I. Fariñas-Sánchez  
Departamento de Física, Universidad Simón Bolívar, Venezuela

Date submitted: 16 Dec 2010

Electronic form version 1.4