

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

Experimental Kataura Plot in Single-walled Carbon Nanotubes

KAIHUI LIU, FAJUN XIAO, RODRIGO CAPAZ, JACK DESLIPPE, Phys Dept. UC Berkeley, WENLONG WANG, IOP, CAS, Beijing, China, SHAUL ALONI, The Molecular Foundry, LBNL, Berkeley, STEVEN LOUIE, Phys Dept. UC Berkeley, ENGE WANG, ICQM, Peking University, Beijing, China, FENG WANG, Phys Dept. UC Berkeley, PHYS DEPT. UC BERKELEY TEAM, THE MOLECULAR FOUNDRY, LBNL, BERKELEY COLLABORATION, IOP, CAS, BEIJING, CHINA COLLABORATION, ICQM, SCHOOL OF PHYSICS, PEKING UNIVERSITY, BEIJING, CHINA COLLABORATION — Single-walled carbon nanotubes (SWNTs) comprise a family of more than 400 structures characterized by different chiral indices n - m , each having a distinct electronic structure that can be either metallic or semiconducting. An outstanding question is how the physical properties, such as optical transitions, vary with the exact nanotube structures. By combining TEM diffraction and Rayleigh scattering spectroscopy on the same individual nanotubes, we determine independently the chiral indices and optical transitions of over 170 single-walled nanotubes. These data permit us to create an experimental Kataura plot for single-walled carbon nanotubes.

Kaihui Liu
Phys Dept. UC Berkeley

Date submitted: 18 Nov 2010

Electronic form version 1.4