

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

Evidence of nodes in the non-centrosymmetric superconductor

Y₂C₃ J. CHEN, Dept. Physics, Zhejiang Univ., H.Q. YUAN, Dept. Physics, Zhejiang Univ., M.B. SALAMON, Dept. Physics, U. Texas at Dallas, S. AKUTAGAWA, J. AKIMITSU, Dept. Physics, Aoyama Gakuin U. — In a non-centrosymmetric superconductor, antisymmetric spin-orbit coupling (ASOC) can admix spin-singlet and spin-triplet pairing states, leading to accidental nodes in the energy gap [1]. Y₂C₃ is such a superconductor with a T_c of 18K. Early NMR and μ SR results [2] indicated a gap structure incompatible with either BCS s-wave or typical d-wave behavior. To further elucidate its superconducting properties, we have measured the temperature dependence of the magnetic penetration depth using a tunneling-diode oscillator technique. While the high temperature penetration depth, and therefore its corresponding superfluid density, can be well described by a two-gap BCS model, as discussed in Ref. [2], the low temperature penetration depth follows a linear temperature dependence, indicating possible existence of nodes in the energy gap. Together with the large upper critical field observed in Y₂C₃ [3], the existence of nodes, we argue, might be attributed to the ASOC as a result of absent inversion symmetry even though other possibilities cannot be excluded. [1] H. Q. Yuan et al, Phys. Rev. Lett. **97**, 017006 (2006). [2] A. Harada et al, J. Phys. Soc. Jpn. **76**, 023704(2007); S. Kuroiwa et al, Phys. Rev. Lett. **100**, 097002 (2008). [3] H. Q. Yuan et al, J. Phys. Chem. Solids (in press).

Myron Salamon
University of Texas at Dallas

Date submitted: 22 Dec 2010

Electronic form version 1.4