

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

Non-Fermi Liquid behavior in Itinerant Ferromagnet, HfMnGa₂

CARLOS MARQUES, Brookhaven National Laboratory and Department of Physics and Astronomy, Stony Brook University, YURI JANSSEN, Brookhaven National Laboratory, MOO SUNG KIM, Brookhaven National Laboratory and Department of Physics and Astronomy, Stony Brook University, LIUSUO WU, Department of Physics and Astronomy, Stony Brook University, MEIGAN ARONSON, Brookhaven National Laboratory and Department of Physics and Astronomy, Stony Brook University — Single crystals of HfMnGa₂, space group Pnma, were grown using a Ga self flux technique. A sharp peak in the AC susceptibility χ_{AC} shows a phase transition at $T_C \approx 26K$, followed by Curie-Weiss behavior at higher temperatures. Arrott plot analysis confirms this transition is ferromagnetic with a spontaneous moment of $\mu_0 \cong 0.3\mu_B/\text{Mn}$. HfMnGa₂ has a coercive field of ~ 0.1 T as well as a large magnetic anisotropy that restricts the moments to point in the [010] direction. Both a large Rhodes-Wohlfarth parameter $\mu_{\text{fluct}}/\mu_0 \cong 3.6$ and low T_C suggest HfMnGa₂ is comparable to other itinerant ferromagnets MnSi, ZrZn₂ and Ni₃Al. Resistivity $\rho(T)$ shows HfMnGa₂ to be metallic with $\rho(T) - \rho_0$ having a $T^{5/3}$ dependence in the ordered state. This non-Fermi liquid relationship was also observed in Ni₃Al and ZrZn₂ over a more limited range of temperatures.

Carlos Marques

Date submitted: 08 Dec 2010

Electronic form version 1.4