

Abstract Submitted  
for the MAR11 Meeting of  
The American Physical Society

**On the Minimum Energy Path to Membrane Pore Formation**

CHRISTINA TING, ZHEN-GANG WANG, California Institute of Technology — Several experimental methods have been developed to study the mechanical response of vesicles under an applied tension. Of particular note are the micropipette aspiration techniques and the use of a viscous solution to extend the lifetime of pores. MD simulations have also been used to study the energetic and structural properties of these transient pores on a molecular level. However, they often require extremely high tensions beyond the regime where pore formation is a thermally-activated event. We approach the nucleation problem by combining the string method with dynamic self-consistent field (DSCF) theory to obtain the full minimum energy path (MEP) to pore formation for a range of surface tensions  $\gamma$ . We compare our results with classical nucleation theory (CNT). Near the coexistence ( $\gamma \rightarrow 0$ ) the rim of the pore is well-defined and the line tension is well approximated by the macroscopic definition given by CNT. However, when the free energy barrier is within  $\sim 10$  kT, the transition state is somewhere between a stalk-like structure and a thinned membrane leading to a hole that is partially exposed to solvents. These molecular rearrangements involved in the formation of a pore are not captured by CNT.

Christina Ting  
California Institute of Technology

Date submitted: 19 Nov 2010

Electronic form version 1.4