

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

Density Functional Theory (DFT) simulations of CO₂ under shock compression and design of liquid CO₂ experiments on Z¹ T. R. MATTSSON, L. SHULENBURGER, S. ROOT, Sandia National Laboratories, Albuquerque, NM., K. R. COCHRANE, Ktech Corporation, Albuquerque, NM. — Quantitative knowledge of the thermo-physical properties of CO₂ at high pressure is required to confidently model the structure of gas-giants like Neptune and Uranus and the deep carbon cycle of the earth. DFT based molecular dynamics has been established as a method capable of yielding high fidelity results for many materials, including shocked gases, at high pressure and temperature. We predict the principal Hugoniot for liquid CO₂ up to 500GPa. Our simulations also show that the plateau in shock pressure identified by Nellis and co-workers [1] is the result of dissociation. At low temperatures we validate the DFT results by comparing with diffusion Monte Carlo calculations. This allows for a more accurate determination of the initial conditions for the shock experiments. We also describe the design of upcoming flyer-plate experiments on the Z-machine aimed at providing high-precision shock compression data for CO₂ between 150 and 600 GPa. [1] W. Nellis, et. al. , J. Chem. Phys. **95**, 5268 (1991).

¹Sandia National Laboratories is a multiprogram laboratory managed and operated by Sandia Corp. for the US Dept. of Energy's National Nuclear Security Administration under Contract DE-AC04-94AL85000.

Thomas R. Mattsson
Sandia National Laboratories

Date submitted: 27 Nov 2010

Electronic form version 1.4