

Abstract Submitted  
for the MAR11 Meeting of  
The American Physical Society

**Measurements of Multiple Gap Substructure in MgB<sub>2</sub> Below 1 Kelvin**<sup>1</sup> STEVEN CARABELLO, JOSEPH LAMBERT, ROBERTO RAMOS, Department of Physics, Drexel University — The two superconducting energy gaps of magnesium diboride (MgB<sub>2</sub>) are well established. First-principles calculations have also predicted substructure within the sigma- and pi-band gaps. However, due to anticipated intraband impurity scattering, there is controversy as to whether these finer structures can be observed in real samples. Prior experimental evidence above 1 Kelvin has provided evidence supportive of these features. We have performed tunneling spectroscopy experiments on MgB<sub>2</sub>/insulator/Pb Josephson junctions on SiC substrates, at temperatures as low as 20 mK. By measuring differential conductance at low temperatures, and by using extremely clean MgB<sub>2</sub> thin films, we have resolved features within the energy gaps to under 1 meV. We report results of these experiments, which are in remarkable agreement with theoretical predictions for this substructure.

<sup>1</sup>We wish to thank Profs. Xiaoxing Xi and Ke Chen for providing high quality MgB<sub>2</sub> Josephson junctions and many helpful insights.

Steven Carabello  
Department of Physics, Drexel University

Date submitted: 19 Nov 2010

Electronic form version 1.4