

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

Raman spectroscopy of graphite in high magnetic fields

YOUNGHEE KIM, National High Magnetic Field Laboratory, ANTONIO LOMBARDO, Engineering Department, Cambridge University, NIKOLAI G. KALUGIN, Department of Materials and Metallurgical Engineering, New Mexico Tech, JUNICHIRO KONO, Department of Electrical & Computer Engineering, Rice University, ANDREA C. FERRARI, Engineering Department, Cambridge University, DMITRY SMIRNOV, National High Magnetic Field Laboratory — Recently, much attention has been paid to electron-phonon coupling in graphene. In particular, significant re-normalization and broadening of long-wavelength optical phonons are predicted to occur through resonant interaction with Landau-quantized Dirac fermions. We report here on a high-field magneto-Raman spectroscopy study of highly-oriented pyrolytic graphite (HOPG) and natural graphite at temperatures down to 5 K and in magnetic fields up to 45 T. The E_{2g} graphite phonon line exhibits anticrossing-like behavior at approximately 30 T, which we attribute to the magneto-phonon resonance (MPR) of graphite's massless holes at the H-point. Additionally, we observed features related to inter-Landau-level transitions at the K-point of graphite. We also observed weak graphene-like signatures of MPR, indicating the existence of graphene flakes on the graphite surface.

Younghee Kim
National High Magnetic Field Laboratory

Date submitted: 28 Nov 2010

Electronic form version 1.4