

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

Bose-Einstein Condensation in Han Purple - a NMR Study¹

RAIVO STERN, NICPB, Tallinn, Estonia, STEFFEN KRÄMER, MLADEN HORVATIC, CLAUDE BERTHIER, LNCMI, Grenoble, France, IVO HEINMAA, ENNO JOON, NICPB, Tallinn, Estonia, TSUYOSHI KIMURA, Osaka University, Japan — NMR study of the two quasi-2D coupled spin-1/2 dimer compound, BaCuSi₂O₆ (Han Purple) [1], is presented. T_{BEC} varies as $(H - H_{c1})^{2/d}$, where d is the dimensionality of the system, and H_{c1} the critical field which closes the gap. BaCuSi₂O₆ was claimed to exhibit an reduction of d from 3D to 2D at low T [2]. However, due to a structural transformation at 90 K, different intradimer exchange couplings and different gaps ($\Delta_B/\Delta_A=1.16$) exist in every second plane along the c axis [3]. In our first NMR experiments [3], we have shown that the population of bosons in the B planes n_B was much smaller than n_A , but finite in the field range $\Delta_A/g\mu_B < H < \Delta_B/g\mu_B$ where $n_B = 0$ is expected in a naive model of uncoupled planes. Recently, a new model has been presented [4] which takes into account both frustration and quantum fluctuations. This leads to a non-zero population n_B of *uncondensed* bosons in the B plane, increasing quadratically with $(H - H_{c1})$, as compared to the linear dependence of n_A . We compare our new NMR results to these predictions. [1] M. Jaime *et al.*, PRL **93**,087203 (2004). [2] S. E. Sebastian *et al.*, Nature **441**, 617 (2006). [3] S. Krämer *et al.*, PRB **76**, 100406(R) (2007). [4] N. Laflorencie and F. Mila, PRL **102**, 060602 (2009).

¹Euromagnet II and Estonian Science Foundation.

Raivo Stern
NICPB

Date submitted: 07 Dec 2010

Electronic form version 1.4