

Abstract Submitted
for the MAR12 Meeting of
The American Physical Society

Sorting Category: 12.1.2 (T)

Enhanced **Opti-**
cal Dichroism of Graphene Nanoribbons¹ VITOR M. PEREIRA,
F. HIPOLITO, Graphene Research Centre and Department of Physics,
National University of Singapore, A.J. CHAVES, R.M. RIBEIRO, M.I.
VASILEVSKIY, N.M.R. PERES, Department of Physics and Centre of
Physics, University of Minho, Portugal. — The optical conductivity of
graphene nanoribbons is analytical and exactly derived. It is shown that
the absence of translational invariance along the transverse direction al-
lows considerable intra-band absorption in a narrow frequency window
that varies with the ribbon width, and lies in the THz band for ribbons
10-100nm wide. In this region the anisotropy in the optical conductivity
can be as high as two orders of magnitude, which renders the medium
dichroic, and allows near 100% polarizability with just a single layer of
graphene. The interplay between the geometrically induced anisotropy
with the anisotropy induced by plasmon absorption is also considered
and discussed.

¹arXiv:1107.1509

Prefer Oral Session
 Prefer Poster Session

Vitor Pereira
vpereira@nus.edu.sg
National University of Singapore

Date submitted: 13 Dec 2011

Electronic form version 1.4