

Abstract Submitted
for the MAR14 Meeting of
The American Physical Society

Hydrogen Trapping in Carbon-Doped h -BN/Rh(111) Nanomesh

JARVIS LOH¹, Michigan Tech Univ, SANDEEP NIGAM, Chemistry Division, Bhabha Atomic Research Centre, Trombay, Mumbai 400085, India, RAVINDRA PANDEY, Department of Physics, Michigan Technological University, Houghton, Michigan 49931, USA, GOVINDA MALLICK, Weapons and Materials Research Directorate, U.S. Army Research Laboratory, Aberdeen Proving Ground, Maryland 21005, USA — Atomic or molecular preferential adsorption on surface nanotemplates provides a facile and feasible means of fabricating ordered low-dimensional nanostructures with tailored functionality. In this study, by employing density-functional theory calculations, we demonstrate (1) the carbon doping of the (B,N)=(fcc_{Rh},top_{Rh}) h -BN/Rh(111) nanomesh, and (2) the selective trapping of hydrogen atoms on these dopants at various sites of the nanomesh – within the pore, on the wire, and at an intermediate site. Contrary to carbon-doped boron nitride sheets, it is energetically more favorable for a carbon impurity to replace a nitrogen atom as compared to a boron atom at all three sites of the nanomesh. In addition, the adsorption energy of hydrogen adsorbates is greater at the wire of a nitrogen-substituted nanomesh relative to that in its pore, while this adsorption energy is invariant at different sites in a boron-substituted nanomesh.

¹Second affiliation: Institute of High Performance Computing, 1 Fusionopolis Way, #16-16 Connexis, Singapore 138632

Jarvis Loh
Michigan Tech Univ

Date submitted: 07 Nov 2013

Electronic form version 1.4