

Abstract Submitted  
for the MAR14 Meeting of  
The American Physical Society

**Shubnikov-de Haas Oscillations in 3D Dirac semimetal Cd<sub>3</sub>As<sub>2</sub>**

SHIH-TING GUO, Institute of Physics, Academia Sinica, Taipei, Taiwan, R. SANKAR, Department of Physics, National Tsing Hua University, Hsinchu, Taiwan, YUNG-YU CHIEN, Institute of Physics, Academia Sinica, Taipei, Taiwan, HORNG-TAY JENG, Center for Condensed Matter Sciences, National Taiwan University, Taipei, Taiwan, F.C. CHOU, Department of Physics, National Tsing Hua University, Hsinchu, Taiwan, WEI-LI LEE, Institute of Physics, Academia Sinica, Taipei, Taiwan, DR.WEI-LI LEE'S LAB, INSTITUTE OF PHYSICS, ACADEMIA SINICA, TAIPEI, TAIWAN TEAM, DEPARTMENT OF PHYSICS, NATIONAL TSING HUA UNIVERSITY, HSINCHU, TAIWAN COLLABORATION, CENTER FOR CONDENSED MATTER SCIENCES, NATIONAL TAIWAN UNIVERSITY, TAIPEI, TAIWAN COLLABORATION — Cadmium arsenide (Cd<sub>3</sub>As<sub>2</sub>) was known for its inverted band structure and ultra-high electron mobility. It has been theoretically predicted to exhibit a 3D Dirac semimetal phase containing degenerate Weyl nodes, which was recently confirmed by ARPES experiments. By conducting magneto-transport measurements in high quality single crystals of Cd<sub>3</sub>As<sub>2</sub>, we found a single frequency  $F = 67$  Tesla in the Shubnikov-de Haas (SdH) oscillations giving an effective mass  $m^* = 0.057 m_e$  with electron mobility as high as 99,500 cm<sup>2</sup>/V-sec, which is consistent with the calculated 3D Dirac semimetal band. In a certain field orientation, we found a splitting of the SdH oscillation frequency in the FFT spectrum, which infers a possible lifting of the degeneracies. Detailed magnetotransport data in several doped and undoped single crystals of Cd<sub>3</sub>As<sub>2</sub> will be presented and discussed.

Shih-Ting Guo  
Academia Sinica

Date submitted: 12 Nov 2013

Electronic form version 1.4