

Abstract Submitted
for the MAR14 Meeting of
The American Physical Society

Spin-orbital separation in the anisotropic ladder system CaCu_2O_3

VALENTINA BISOGNI, Paul Scherrer Institut, Villigen PSI, Switzerland, KRZYSZTOF WOHLFELD, SLAC National Laboratory and Stanford University, Menlo Park, USA, SATOSHI NISHIMOTO, IFW Dresden, Germany, CLAUDE MONNEY, Fritz-Haber-Insitut, Berlin, Germany, JAN TRINCKAUF, IFW Dresden, Germany, KEJIN ZHOU, Diamond Light Source, Didcot, UK, ROBERTO KRAUS, IFW Dresden, Germany, THORSTEN SCHMITT, Paul Scherrer Institut, Villigen PSI, Switzerland, JEROEN VAN DEN BRINK, JOCHEN GECK, IFW Dresden, Germany — Recently, resonant inelastic X-ray scattering (RIXS) on the 1D spin system Sr_2CuO_3 has revealed an unprecedented dispersion of orbital excitations [Nature 485, 82 (2012)]. This result has been interpreted as the fractionalization of spin and orbital degree of freedom from the elementary electron, hallmark of one dimensional physics as the previously observed spin-charge separation [Nature Phys. 2, 397 (2006)]. How these phenomena carry over into higher dimensions remains currently unclear. To clarify this point, we studied the spin and orbital excitations of the anisotropic ladder CaCu_2O_3 , which realizes a first step towards 2D correlated electron systems. Combining high-resolution RIXS experiments with theoretical model calculations we show that spin-orbital fractionalization indeed occurs in CaCu_2O_3 and prevails beyond the strict 1D limit [arXiv:1310.8346]. We also establish that such a fractionalization is far more robust than the spin-charge separation. The main reasons behind this are the intrinsic 1D orbital dynamics and the fact that the spinons are faster than the orbitons but slower than the holons.

Thorsten Schmitt
Paul Scherrer Institut, Villigen PSI, Switzerland

Date submitted: 15 Nov 2013

Electronic form version 1.4