

Abstract Submitted  
for the MAR14 Meeting of  
The American Physical Society

**Exceptional ballistic transport in epitaxial graphene nanoribbons**

CHRISTOPH TEGENKAMP, JENS BARINGHAUS, FREDERIK EDLER, Institut für Festkörperphysik, Leibniz Universität Hannover, CLAIRE BERGER, WALT A. DE HEER, School of Physics, Georgia Institute of Technology, Atlanta, USA — The patterning of graphene into graphene nanoribbons is an essential task for the development of graphene based devices. For such ribbons with a well-ordered edge geometry the presence of one-dimensional edge states has been predicted. We use a selective graphitization process on SiC-mesa structures to produce graphene nanoribbons with a width of 40 nm. The local electronic properties of the ribbons are investigated by means of a 4-tip STM. In combination with a SEM, the precise positioning of all four tips on the nanometer range is possible to perform local transport measurements. Furthermore, local tunneling spectroscopy reveals characteristic features of ferromagnetic zig-zag graphene nanoribbons. Transport experiments carried out on the very same ribbon show a conductance of  $e^2/h$  for a wide temperature range from 30 K up to room temperature and probe spacings between 1  $\mu m$  and 10  $\mu m$ . Description within the Landauer formalism is possible assuming ballistic transport dominated by a single channel. Transport in the second zeroth subband is only detectable for probe spacings smaller than 1  $\mu m$  due to the short localization length of carriers in this subband manifesting in the increase of the conductance to  $2e^2/h$  at probe spacings below 200 nm.

Christoph Tegenkamp  
Institut für Festkörperphysik, Leibniz Universität Hannover

Date submitted: 15 Nov 2013

Electronic form version 1.4