

Abstract Submitted  
for the MAR14 Meeting of  
The American Physical Society

**Weak phase stiffness and nature of the quantum critical point in underdoped cuprates**<sup>1</sup> WEI KU, YUCEL YILDIRIM<sup>2</sup>, Brookhaven National Laboratory — We demonstrate that the zero-temperature superconducting phase diagram of underdoped cuprates can be quantitatively understood in the strong binding limit, using only the experimental spectral function of the “normal” pseudo-gap phase without any free parameter. In the prototypical  $(\text{La}_{1-x}\text{Sr}_x)_2\text{CuO}_4$ , a kinetics-driven  $d$ -wave superconductivity is obtained above the critical doping  $\delta_c \sim 5.2\%$ , below which complete loss of superfluidity results from local quantum fluctuation involving local  $p$ -wave pairs. Near the critical doping, a enormous mass enhancement of the local pairs is found responsible for the observed rapid decrease of phase stiffness. Finally, a striking mass divergence is predicted at  $\delta_c$  that dictates the occurrence of the observed quantum critical point and the sudden suppression of the Nernst effects in the nearby region.

<sup>1</sup>Work supported by U.S. Department of Energy, Office of Basic Energy Science, under Contract No. DE-AC02-98CH10886

<sup>2</sup>current address: Physics Department, Faculty of Arts and Sciences, Doğuş University, Acıbadem-Kadıköy, 34722 Istanbul, Turkey

Wei Ku  
Brookhaven National Laboratory

Date submitted: 15 Nov 2013

Electronic form version 1.4