

Abstract Submitted
for the MAR15 Meeting of
The American Physical Society

Strong-randomness

phenomena in quantum Ashkin-Teller models¹ THOMAS VOJTA, HATEM BARGHATHI, FAWAZ HRAHSHEH, Missouri Univ of Sci & Tech, JOSE HOYOS, Instituto de Fisica de Sao Carlos, Universidade de Sao Paulo, RAJ NARAYANAN, Indian Institute of Technology Madras — The N -color quantum Ashkin-Teller spin chain is a prototypical model for the study of strong-randomness phenomena at first-order and continuous quantum phase transitions. This talk discusses strong-disorder renormalization group approaches to this system in the weak-coupling as well as the strong-coupling regimes. Specifically, we introduce a novel general variable transformation that unifies the treatment of the strong-coupling regime. This allows us to determine the phase diagram for all color numbers N , and the critical behavior for all $N \neq 4$. In the case of two colors, $N = 2$, a partially ordered product phase separates the paramagnetic and ferromagnetic phases in the strong-coupling regime. This phase is absent for all $N > 2$, i.e., there is a direct phase boundary between the paramagnetic and ferromagnetic phases. In agreement with the quantum version of the Aizenman-Wehr theorem, all phase transitions are continuous, even if their clean counterparts are of first order. We also discuss the various critical and multicritical points. They are all of infinite-randomness type, but depending on the coupling strength, they belong to different universality classes.

¹We are grateful for the support from NSF under Grant Nos. DMR-1205803 and PHYS-1066293, from Simons Foundation, from FAPESP under Grant No. 2013/09850-7, and from CNPq under Grant Nos. 590093/2011-8 and 305261/2012-6.

Thomas Vojta
Missouri Univ of Sci & Tech

Date submitted: 04 Nov 2014

Electronic form version 1.4