

Abstract Submitted
for the MAR15 Meeting of
The American Physical Society

Reflective Optical Limiter Based on Resonant Transmission¹

ELEANA MAKRI, TSAMPIKOS KOTTOS, Department of Physics, Wesleyan University, Middletown CT-06459, USA, ILYA VITEBSKIY, Air Force Research Laboratory, Sensors Directorate, Wright Patterson AFB, OH 45433 USA — Optical limiters transmit low-level radiation while blocking electromagnetic pulses with excessively high energy (energy limiters) or with excessively high peak intensity (power limiters). A typical optical limiter absorbs most of the high-level radiation which can cause its overheating and destruction. Here we introduce the novel concept of a reflective energy limiter which blocks electromagnetic pulses with excessively high total energy by reflecting them back to space, rather than absorbing them. The idea is to use a defect layer with temperature dependent loss tangent embedded in a low-loss photonic structure. The low energy pulses with central frequency close to that of the localized defect mode will pass through. But if the cumulative energy carried by the pulse exceeds certain level, the entire photonic structure becomes highly reflective (not absorptive!) within a broad frequency range. The underlying physical mechanism is based on self-regulated impedance mismatch which increases dramatically with the cumulative energy carried by the pulse.

¹AFOSR MURI FA9550-14-1-0037

Eleana Makri
Wesleyan University

Date submitted: 04 Nov 2014

Electronic form version 1.4