

Abstract Submitted
for the MAR15 Meeting of
The American Physical Society

Superfluidity in Strongly Interacting Spin-Polarized Fermi Gases¹

BEN A. OLSEN, MELISSA REVELLE, JACOB A. FRY, RANDALL G. HULET, Department of Physics & Astronomy and Rice Quantum Institute, Rice University, Houston, TX 77005, DANIEL E. SHEEHY, Department of Physics & Astronomy, Louisiana State University, Baton Rouge, LA 70803 — We report measurements of phase separation in a harmonically trapped, spin polarized two-component Fermi gas. The interactions in the gas are varied using a magnetically-tuned Feshbach resonance between the weakly-interacting BCS and strongly-interacting BEC regimes. Using spin-selective imaging, we measure the density profiles for the two lowest hyperfine levels of ^6Li , with the superfluid phase being indicated by an unpolarized central core. We determine phase boundaries for the superfluid transition as a function of interactions and polarization, finding results that are consistent with earlier experimental results in the crossover regime². We also explore the BEC side of resonance, where we compare our measurements to Quantum Monte Carlo simulations³, and also the deep BCS regime, where few theoretical predictions are available.

¹Supported by DARPA, NSF, ARO, and ONR

²M. Zwierlein et al., *Science* **311**, 492 (2006); N. Navon et al., *Science* **328**, 729 (2010)

³G. Bertaina and S. Giorgini, *PRA* **79**, 013616 (2009)

Ben Olsen
Rice University

Date submitted: 14 Nov 2014

Electronic form version 1.4