

Abstract Submitted
for the MAR15 Meeting of
The American Physical Society

Current flow in biased bilayer graphene: the role of sublattices

CARLOS PAEZ, Universidade Estadual de Campinas-FCA, DARIO BAHAMON, MackGraphe -Graphene and Nano-Materials Research Center, Mackenzie Presbyterian University, ANA PEREIRA, Universidade Estadual de Campinas-FCA — We investigate here how the current flows over a bilayer graphene in the presence of an external electric field perpendicularly applied (biased bilayer). Charge density polarization between layers in these systems is known to create a layer pseudospin, which can be manipulated by the electric field. Our results show that current does not necessarily flow over regions of the system with higher charge density. Charge can be predominantly concentrated over one layer, while current flows over the other layer. We find that this phenomenon occurs when the charge density becomes highly concentrated over only one of the sublattices, as the electric field breaks layer and sublattice symmetries for a Bernal-stacked bilayer. For bilayer nanoribbons, the situation is even more complex, with a competition between edge and bulk effects for the definition of the current flow. We show that, in spite of not flowing through the layer where charge is polarized to, the current in these systems also defines a controllable layer pseudospin.

Carlos Paez
Universidade Estadual de Campinas-FCA

Date submitted: 24 Nov 2014

Electronic form version 1.4