

Abstract Submitted
for the MAR16 Meeting of
The American Physical Society

Unliganded EphA3 dimerization promoted by the SAM domain

KALINA HRISTOVA, DEO SINGH, CHRISTOPHER KING, FOZIA AHMED, Johns Hopkins University, ELENA PASQUALE, Sanford-Burnham Medical Research Institute — The EphA3 receptor tyrosine kinase regulates morphogenesis during development, and is overexpressed and mutated in a variety of cancers. EphA3 activation is believed to follow a “seeding mechanism” model, in which ligand binding to the monomeric receptor acts as a trigger for signal-productive receptor clustering. We use a novel approach to study EphA3 lateral interactions on the surface of live cells, and we demonstrate that EphA3 forms dimers in the absence of ligand binding. We further show that these dimers are stabilized by interactions involving the EphA3 SAM domain. The discovery of unliganded EphA3 dimers challenges the current understanding of the chain of EphA3 activation events, and suggests that EphA3 may follow the “pre-formed dimer” model of activation known to be relevant for other receptor tyrosine kinases. This work also establishes a new role for the SAM domain in promoting Eph receptor lateral interactions and signaling on the cell surface.

Kalina Hristova
Johns Hopkins University

Date submitted: 29 Oct 2015

Electronic form version 1.4