

Abstract Submitted
for the MAR16 Meeting of
The American Physical Society

Spontaneous symmetry breaking in correlated wave functions

RYUI KANEKO, Goethe-Universitaet Frankfurt, LUCA F. TOCCHIO, Consiglio Nazionale delle Ricerche (CNR) and International School for Advanced Studies (SISSA), ROSER VALENTI, Goethe-Universitaet Frankfurt, FEDERICO BECCA, Consiglio Nazionale delle Ricerche (CNR) and International School for Advanced Studies (SISSA), CLAUDIUS GROS, Goethe-Universitaet Frankfurt — We show that Jastrow-Slater wave functions, in which a density-density Jastrow factor is applied onto an uncorrelated fermionic state, may possess long-range order even when all symmetries are preserved in the wave function. This fact is mainly related to the presence of a sufficiently strong Jastrow term (also including the case of full Gutzwiller projection, suitable for describing spin models). Selected examples are reported, including the spawning of Néel order and dimerization in spin systems, and the stabilization of density and orbital order in itinerant electronic systems. [1] Ryui Kaneko, Luca F. Tocchio, Roser Valentí, Federico Becca, and Claudius Gros, arXiv:1510.08653.

Ryui Kaneko
Goethe-Universitaet Frankfurt

Date submitted: 05 Nov 2015

Electronic form version 1.4