

Abstract Submitted
for the MAR16 Meeting of
The American Physical Society

Probing moiré physics in low angle twisted bilayer graphene

YUAN CAO, JASON LUO, Massachusetts Inst of Tech-MIT, JAVIER D. SANCHEZ-YAMAGISHI, Harvard University, KENJI WATANABE, TAKASHI TANIGUCHI, NIMS, Japan, PABLO JARILLO-HERRERO, Massachusetts Inst of Tech-MIT — When two sheets of honeycomb structured graphene are stacked together, a moiré pattern that depend on the twist angle forms. A gapped superlattice band structure is resulted when this angle is small enough so that the energy of moiré modulated hybridization between wave functions on the two sheets that are separated by only 0.34nm is low enough. Apart from the energy gaps, the superlattice band structure also manifests reduced Fermi velocity which implies comparatively higher density of states, multiple van Hove singularities below and above the gaps, and Hofstadter butterfly physics when magnetic field is applied. We show electronic transport measurements of high-quality low-angle twist bilayer graphene devices fabricated by a novel tear-and-stack technique, at zero, low and high magnetic fields. We also present angle dependence of the electronic structure along with magneto-transport features that possibly imply electron-electron interactions. A brief discussion about the transition between the low-twist and high-twist bilayer graphene, the physics of the latter of which is believed to be essentially based on decoupled monolayer graphene according to our previous work, is included.

Yuan Cao
Massachusetts Inst of Tech-MIT

Date submitted: 07 Nov 2015

Electronic form version 1.4