

Abstract Submitted  
for the MAR16 Meeting of  
The American Physical Society

**Mott Quantum Criticality in the Anisotropic 2D Hubbard Model<sup>1</sup>**

BENJAMIN LENZ, SALVATORE R. MANMANA, THOMAS PRUSCHKE, Institute for Theoretical Physics, University of Göttingen, FAKHER F. ASSAAD, MARCIN RACZKOWSKI, Institute for Theoretical Physics and Astrophysics, University of Würzburg — We present evidence for Mott quantum criticality in an anisotropic two-dimensional system of coupled Hubbard chains at half-filling. In this scenario emerging from variational cluster approximation and cluster dynamical mean-field theory, the interchain hopping  $t_{\perp}$  acts as control parameter driving the second-order critical endpoint  $T_c$  of the metal-insulator transition down to zero at  $t_{\perp}^c/t \simeq 0.2$ . Below  $t_{\perp}^c$  the volume of hole and electron Fermi pockets of a compensated metal vanishes continuously at the Mott transition. Above  $t_{\perp}^c$  the volume reduction of the pockets is cut off by a first-order transition. We discuss the relevance of our findings to a putative quantum critical point in layered organic conductors whose location remains elusive so far.

<sup>1</sup>We acknowledge support by DFG research units FOR1807 and FOR1346, ERC Starting Grant No. 306897 and NSF Grant No. PHY11-25915, and computer support by the GWDG and Jülich Supercomputing Centre.

Benjamin Lenz  
Institute for Theoretical Physics, University of Göttingen

Date submitted: 06 Nov 2015

Electronic form version 1.4