

Abstract Submitted  
for the MAR16 Meeting of  
The American Physical Society

**Magnetic Correlations in the Triangular Antiferromagnet TbInO<sub>3</sub>**

GABRIELE SALA, McMaster University, LUCY CLARK, University of St Andrews, DALINI MAHARAJ, McMaster University, MATTHEW B. STONE, Oak Ridge National Laboratory, KEVIN S. KNIGHT, ISIS Neutron Facility, SANGWOOK CHEONG, Rutgers University, BRUCE D. GAULIN, McMaster University — TbInO<sub>3</sub> crystallizes with a hexagonal  $P6_3cm$  structure in which layers of edge-sharing triangles of magnetic Tb<sup>3+</sup> ions are separated by non-magnetic [InO<sub>5</sub>]<sup>7-</sup> units. TbInO<sub>3</sub>, therefore, realizes an excellent opportunity to explore the behavior of a two-dimensional magnetic triangular lattice, a canonical model of geometric frustration. Here we present our study of a polycrystalline sample of TbInO<sub>3</sub>. Our high resolution powder neutron diffraction data (HRPD, ISIS) of TbInO<sub>3</sub> confirm that the triangular layers of Tb<sup>3+</sup> remain undistorted to at least 0.46 K. Magnetic susceptibility data follow Curie-Weiss behavior over a wide range of  $T$  with  $\theta = -17.19(3)$  K indicating the dominance of antiferromagnetic correlations. The susceptibility data also show an absence of conventional long-range spin order down to at least 0.55 K, reflecting the frustrated nature of TbInO<sub>3</sub>. Elastic magnetic diffuse neutron scattering (SEQUOIA, SNS) is observed below  $\sim 15$  K, due to the presence of static two-dimensional spin correlations. The spectrum of crystal field excitations in TbInO<sub>3</sub> appears to have an exotic form due to the existence of two crystallographically distinct Tb<sup>3+</sup> sites and leads to a strong Ising anisotropy of the spin symmetry.

Lucy Clark  
University of St Andrews

Date submitted: 06 Nov 2015

Electronic form version 1.4