

Abstract Submitted
for the MAR16 Meeting of
The American Physical Society

Interaction of Molecular Oxygen with Au (001) Surface¹

MICHAEL PIERCE, ANDREW LOHEAC², Rochester Institute of Technology, ANDI BARBOUR, Brookhaven National Laboratory, VLADIMIR KOMANICKY, Safarik University, ANTHONY RUFFINO, Rochester Institute of Technology, HOYDOO YOU, Argonne National Laboratory — Kinetics of dioxygen - Au (001) surface interaction has been studied at high temperature and near atmospheric pressures with in situ x-ray scattering measurements. We find that the hexagonal reconstruction (hex) of Au (001) surface lifts to (11) in the presence of dioxygen. The measured lifting rate constant vs. temperature shows a ‘volcano’-type behavior indicating that oxygen adsorption limits at high temperature while activation barrier of the lifting limits at low temperature. The pressure–temperature (PT) phase diagram obtained in our study shows three regions: hex at low P and T, (11) at high P and T, and coexistence of the hex and the (11) at the intermediate P and T.

¹Funding provided by Research Corporation for Science Advancement. Work done at the Advanced Photon Source supported by the U.S. Department of Energy.

^{2*} Now at University of North Carolina - Chapel Hill

Michael Pierce
Rochester Institute of Technology

Date submitted: 06 Nov 2015

Electronic form version 1.4