

Abstract Submitted
for the MAR17 Meeting of
The American Physical Society

Fundamentals of CO₂ adsorption on model copper oxide surfaces

STEVEN J. TJUNG, JACOB J. REPICKY, NANCY M. SANTAGATA, XIAOWA NIE, SIMUCK F. YUK, ARAVIND ASTHAGIRI, JAY A. GUPTA, The Ohio State University — Copper oxides can catalyze the conversion of CO₂ selectively into methanol. We are studying fundamental aspects of the adsorption of CO₂ on model copper oxide surfaces using low temperature (5 K) scanning tunneling microscopy (STM) and spectroscopy (STS). We prepared an oxygen adlayer by exposing Cu(100) to O₂ at 575K, which results in a $(2\sqrt{2}\times\sqrt{2})R45^\circ$ missing row reconstruction. STM images of the pristine surface show a ladder-like contrast with distinct domains. STS reveals a series of image potential states consistent with an increase in work function and two additional unoccupied states. We introduce CO₂ to the surface in situ, which allows us to unambiguously identify individual CO₂ molecules and their adsorption sites. We find that the CO₂ molecules sit in between the O atoms in the missing row reconstruction. The CO₂ molecules are easily perturbed by the STM tip under typical imaging conditions, suggesting that the molecules are weakly bound to the surface. STS is used to probe molecular orbital states of CO₂ molecules. Density functional theory calculations of adsorption sites, vibrational modes, and diffusion barriers are in qualitative agreement with the experiment.

Steven Tjung
Ohio State Univ - Columbus

Date submitted: 11 Nov 2016

Electronic form version 1.4