

Abstract Submitted  
for the MAR17 Meeting of  
The American Physical Society

**Temperature and Magnetic Field Effects on the Raman Spectra of TaSe<sub>2</sub>** JACOB HARDING, J.R. SIMPSON, Towson University, A.R. HIGHT WALKER, NIST — In bulk form, TaSe<sub>2</sub> exhibits transitions between commensurate and incommensurate charge-density wave (CDW) phases, and is attracting interest for advance device applications. In order to explore the evolution of the groundstate CDW phase, mechanical exfoliation of bulk crystals provides freshly cleaved surfaces and may be used to prepare few- to single-layer flakes. In the present work, we extended our opto-thermal Raman measurements [1] on MoS<sub>2</sub> to include other TMDs, specifically TaSe<sub>2</sub>, in both *1T* and *2H* crystallographic phases. A novel, magneto-Raman microscope system affords measurement of low-frequency (down to 10cm<sup>-1</sup>) vibrational modes as a function of both temperature (~10K to 300K) and magnetic field (0T to 9T). The dependence of the observed Raman-active phonons on temperature and magnetic field will be discussed and compared with earlier results on MoS<sub>2</sub>. Specifically, we observe the appearance of low-frequency, zone-folded modes in the CDW state, which soften with temperature similar to the higher frequency, in-plane *E*<sub>2g</sub> mode. Additionally, magnetic-field dependence, including Faraday rotation in the micro-crystal insert will be discussed. [1] R. Yan, J. R. Simpson, et al., ACS Nano 8, 986 (2014).

Jacob Harding  
Towson University

Date submitted: 11 Nov 2016

Electronic form version 1.4