

SES17-2017-000083

E

Abstract for an Invited Paper
for the SES17 Meeting of
the American Physical Society

Current Results of the PRad Experiment at JLab¹

MAXIME LEVILLAIN, North Carolina AT State University, PRAD COLLABORATION

The latest measurements of the proton radius through muonic hydrogen Lamb shift show a discrepancy of about 7σ from a global analysis of standard hydrogen Lamb shift and elastic ep -scattering. In order to understand this *proton radius puzzle*, the PRad experiment, that was designed through an independent method, successfully performed in June 2016, taking elastic ep and Møller-scattering data with rich statistics in a wide Q^2 range including very low momentum transfer ($Q^2 \in [10^{-4}, 10^{-1}] \text{ GeV}^2$) with very accurate angle and energy measurements to minimize the systematic uncertainties to achieve a sub-percent precision. After briefly reminding the framework around this *proton radius puzzle* and how the specific setup of the PRad experiment was designed to improve the previous measurements on elastic ep -scattering cross-sections, using a high efficiency and high resolution calorimeter (HyCal) and a high resolution GEM detector, the presentation will focus on the data analysis and the extraction of ep scattering cross-section normalized by the well known Møller cross-section, to finally show the latest physics results from the 2.2 GeV data.

¹The PRad experiment is supported in part by NSF MRI award PHY-1229153