

Abstract Submitted  
for the TSF16 Meeting of  
The American Physical Society

**Coherent Diffractive Imaging in the Near Field at Large Angles**

BENJAMIN A. POUND, KIMBERLEY NGUYEN, MATTHEW C. TYSON, JOHN L. BARBER, RICHARD L. SANDBERG, Los Alamos National Laboratory — Coherent diffraction imaging (CDI) is a rapidly developing form of lensless imaging where the intensity of the diffraction pattern is directly imaged on a CCD and iterative phase retrieval (IPR) algorithms are used to reconstruct a high resolution image of the sample. This is especially useful at x-ray wavelengths, where lenses are inefficient and difficult to manufacture. However, one challenge with CDI is that the exact relationship between light emerging from the sample and arriving at the detector is nearly impossible to determine, even numerically, but becomes tractable with various assumptions. The standard far-field assumptions require the detector to be placed hundreds or thousands of meters from the sample at hard x-ray energies, which is not practical. A different set of assumptions called the distorted object approach allows imaging at any distance, but has the strict requirement that a value called the small angle number,  $An$ , needs to be much smaller than one. Here we examine where the distorted object approach fails, specifically in regards to  $An$ . We have found that we can obtain good quality images with very large  $An$  values at visible wavelengths.

Richard L. Sandberg  
Los Alamos National Laboratory

Date submitted: 30 Sep 2016

Electronic form version 1.4