

Abstract Submitted
for the TSS18 Meeting of
The American Physical Society

Two-body scattering resonances in an effectively one-dimensional Rashba-Dresselhaus spin-orbit coupled quantum gas¹ SU-JU WANG, QINGZE GUAN, D. BLUME, University of Oklahoma — Confinement-induced resonances allow for the tuning of the effective one-dimensional coupling constant. When the scattering state associated with the ground transverse mode is brought into resonance with the bound state attached to the energetically excited transverse modes, the atoms interact through an infinitely strong repulsion. This provides a route to realize the Tonks-Girardeau gas. On the other hand, the realization of synthetic gauge fields in cold atomic systems has attracted a lot of attention. For instance, bound-state formation is found to be significantly modified in the presence of spin-orbit coupling in three dimensions. This motivates us to study ultracold collisions between two Rashba-Dresselhaus spin-orbit coupled atoms in a quasi-one-dimensional geometry. We develop a multi-channel scattering formalism that accounts for the external transverse confinement and the spin-orbit coupling terms. The interplay between these two single-particle terms is shown to give rise to new scattering resonances. In particular, we focus our discussions on the case with a singlet two-body interaction.

¹Support by the NSF is gratefully acknowledged.

Su-Ju Wang
University of Oklahoma

Date submitted: 09 Mar 2018

Electronic form version 1.4