

Abstract Submitted
for the APR17 Meeting of
The American Physical Society

Resonant tidal excitation of superfluid neutron stars in coalescing binaries HANG YU, NEVIN WEINBERG, Massachusetts Inst of Tech-MIT — We study the resonant tidal excitation of g-modes in coalescing superfluid neutron star (NS) binaries and investigate how such tidal driving impacts the gravitational-wave signal of the inspiral. Previous studies treated the NS core as a normal fluid and did not account for its superfluidity. The source of buoyancy that supports the g-modes is fundamentally different in the two cases: in a normal fluid core the buoyancy is due to gradients in the proton-to-neutron fraction whereas in a superfluid core it is due to gradients in the muon-to-electron (or hyperon) fraction. The latter yields a stronger stratification and a superfluid NS has a denser spectrum of g-modes. As a result, many more g-modes undergo resonant tidal excitation during the inspiral. We find that $\simeq 10$ times more orbital energy is transferred into g-mode oscillations if the NS has a superfluid core rather than a normal fluid core. However, because this energy is transferred later in the inspiral when the orbital decay is faster, the accumulated phase error in the gravitational waveform is comparable for a superfluid and normal fluid NS ($\sim 10^{-3} - 10^{-2}$ rad). A phase error of this magnitude is too small to be measured with the current generation of gravitational wave detectors.

Hang Yu
Massachusetts Inst of Tech-MIT

Date submitted: 25 Sep 2016

Electronic form version 1.4