

Abstract Submitted
for the DAMOP10 Meeting of
The American Physical Society

Optical transition of the ^{229}Th nucleus in a solid-state environment¹ WADE RELLERGERT, SCOTT SULLIVAN, RADOYEH SHOJAEI, UCLA, DAVID DEMILLE, Yale, RICHARD GRECO, MARKUS HEHLEN, JUSTIN TORGERSON, LANL, ERIC HUDSON, UCLA, THE THOR COLLABORATION — We describe a novel approach to directly measure the energy of the narrow, low-lying isomeric state in ^{229}Th . Since nuclear transitions are far less sensitive to environmental conditions than atomic transitions, we argue that the ^{229}Th optical nuclear transition may be driven inside a host crystal with a high transition Q. This technique might also allow for the construction of a solid-state optical frequency reference that surpasses the precision of current optical clocks, as well as improved limits on the variability of fundamental constants. Based on analysis of the crystal lattice environment, we argue that a precision of $3 \cdot 10^{-17} < \Delta f/f < 1 \cdot 10^{-15}$ after 1 s of photon collection may be achieved with a systematic-limited accuracy of $\Delta f/f \sim 2 \cdot 10^{-16}$. Improvement by a factor of 10^2 to 10^3 of the constraints on the variability of several important fundamental constants also appears possible. We report on progress towards evaluation of candidate host crystals.

¹We acknowledge funding from UCOP LRP program.

Scott Sullivan
UCLA

Date submitted: 22 Jan 2010

Electronic form version 1.4