

Abstract Submitted
for the DAMOP16 Meeting of
The American Physical Society

Double Photoionization of Beryllium atoms using Effective Charge approximation HARIPADA SAHA, University of Central Florida, Orlando, FL 32816 — We plan to report the results of our investigation on double photoionization K-shell electrons from Beryllium atoms. We will present the results of triple differential cross sections at excess energy of 20 eV using our recently extended MCHF method [1]. We will use multiconfiguration Hartree Fock method to calculate the wave functions for the initial state. The final state wave functions will be obtained in the angle depended Effective Charge approximation [2-4] which accounts for electron correlation between the two final state continuum electrons. We will discuss the effect of core correlation and the valence shell electrons in the triple differential cross section. The results will be compared with the available accurate theoretical calculations and experimental findings. [1] Hari P. Saha, Phys. Rev. A 87, 042703 (2013). [2] M.R.H. Rudge, Rev. Mod. Phys. 40, 564 (1968). [3] D. Proulx and R. Shakeshaft,, Phys. Rev A 48, R875 (1993). [4]; M. Pont and R. Shakeshaft, Phys. Rev. A51, R2676 (1995).

Haripada Saha
University of Central Florida, Orlando, FL 32816

Date submitted: 08 Jan 2016

Electronic form version 1.4