

Abstract Submitted
for the MAR08 Meeting of
The American Physical Society

Molecular dynamics simulations of hexane on graphite at various coverages: the difference explicit hydrogens make¹ M.W. ROTH, M.J. CONNOLLY, University of Northern Iowa, Department of Physics, PAUL A. GRAY, University of Northern Iowa, Department of Computer Science, CARLOS WEXLER, University of Missouri Columbia, Department of Physics and Astronomy — Molecular Dynamics simulations of hexane (C_6H_{14}) adlayers on graphite are carried out for coverages of $0.5 \leq \rho \leq 1$ monolayers. The hexanes have explicit hydrogens and the graphite is modeled as an all-atom, six-layer structure. Above $\rho \cong 0.9$ the herringbone solid loses orientational order at $T_1 = 140 \text{ K} \pm 3 \text{ K}$. At $\rho = 0.878$ the system presents vacancy patches and T_1 decreases to ca. 100 K. As coverage decreases further, the vacancy patches become larger and by $\rho = 0.614$ the solid is a connected network of randomly oriented domains. All cases show a weak nematic mesophase. The melting temperature is $T_2 = 160 \text{ K} \pm 3 \text{ K}$ and falls to ca. 145 K by $\rho = 0.614$. The dynamics and energetics observed demonstrate that the explicit-hydrogen model of hexane is substantially more realistic than the UA approximation.

¹Acknowledgment is made to the Donors of The American Chemical Society Petroleum Research Fund (PRF43277 - B5), the University of Missouri Research Board, and the D.O.E. (DE-FG02-07ER46411).

Michael Roth
University of Northern Iowa

Date submitted: 12 Dec 2007

Electronic form version 1.4