

Abstract Submitted  
for the MAR08 Meeting of  
The American Physical Society

**Squeezed State Effects on Continuous Variable Quantum Erasing**

PETER BONANNO, VIJAY KASISOMAYAJULA, ONOFRIO RUSSO, New Jersey Institute of Technology — Experimental verification of complementarity using quantum erasing for the continuous variable (CV) infinite dimensional Hilbert space has been considered. [1] The complementary pair is that of the canonically conjugate amplitude and phase quadratures of light. The amplitude quadrature is labeled to a squeezed meter signal by quantum nondemolition (QND) [2] entanglement coupling. [3] Knowledge of *which eigenstate* (WE) can be obtained by measuring this amplitude in the meter state, and can thereafter be ‘lost’ by measuring the quadrature phase of the meter, thus restoring the quadrature phase of the signal beam in a process known as *quantum erasure*. [4] The coupling, i.e. the labeling of the signal state to the meter state, is implemented with a beam splitter coupled to the squeezed light meter beam. [4] We investigate the effects of using the unitary squeeze operator  $S(z)=\exp\frac{1}{2}(z^*a^2 - za^{+2})$  where  $z = re^{i(\text{squeezingangle})}$  on selected coherent states under certain conditions. [5,6] [1] U. L. Anderson et al., Phys. Rev. Lett. 93, 100403 (2004). [2] V. B. Braginsky et al., Science 209, 547 (1980). [3] R. Bruckmeier et al., Phys. Rev. Lett. 79, 43 (1997). [4] P. Grangier et al., Nature 396, 537 (1998). [5] C. M. Caves, Phys. Rev. D 23, 1693 (1981). [6] D. Stoler, Phys. Rev. D. 1, 3217 (1970), D. Stoler, Phys. Rev. D. 4, 1925 (1971). .

Onofri Russo  
New Jersey Institute of Technology

Date submitted: 05 Dec 2007

Electronic form version 1.4