

Abstract Submitted  
for the MAR17 Meeting of  
The American Physical Society

**Scanning tunneling spectroscopy as a probe of multi- $\mathbf{Q}$  magnetic states of itinerant magnets** MARIA NAVARRO GASTIASORO, Niels Bohr Inst, ILYA EREMIN, Institut fr Theoretische Physik III, Ruhr-Universitt Bochum, 44801 Bochum, Germany, RAFAEL M FERNANDES, School of Physics and Astronomy, University of Minnesota, Minneapolis, Minnesota 55455, USA, BRIAN M ANDERSEN, Niels Bohr Inst — The combination of electronic correlations and Fermi surfaces with multiple nesting vectors can lead to the appearance of complex multi- $\mathbf{Q}$  magnetic ground states, hosting unusual states such as chiral density-waves and quantum Hall insulators. Distinguishing single- $\mathbf{Q}$  and multi- $\mathbf{Q}$  magnetic phases is however a notoriously difficult experimental problem. Here we propose theoretically that the local density of states (LDOS) near a magnetic impurity, whose orientation may be controlled by an external magnetic field, can be used to map out the detailed magnetic configuration of an itinerant system and distinguish unambiguously between single- $\mathbf{Q}$  and multi- $\mathbf{Q}$  phases. We demonstrate this concept by computing and contrasting the LDOS near a magnetic impurity embedded in three different magnetic ground states relevant to iron-based superconductors – one single- $\mathbf{Q}$  and two double- $\mathbf{Q}$  phases. Our results open a promising avenue to investigate complex magnetic configurations in itinerant systems via standard scanning tunneling spectroscopy, without requiring spin-resolved capability.

Maria Navarro Gastiasoro  
Niels Bohr Inst

Date submitted: 10 Nov 2016

Electronic form version 1.4