

Abstract Submitted
for the MAR17 Meeting of
The American Physical Society

Universal Dynamics of a localized excitation after an interaction quench¹ FABIO FRANCHINI, Institut Ruer Bokovi, FABIO FRANCHINI TEAM, MANAS KULKARNI TEAM, ANDREY GROMOV TEAM, ANDREA TROMBETTONI TEAM — We study the time evolution -induced by a quench- of local excitations in one dimension. We focus on interaction quenches: the considered protocol consists in creating a stable localized excitation propagating through the system, and then operating a sudden change of the interaction between the particles. To highlight the effect of the quench, we take the initial excitation to be a soliton. The quench splits the excitation into two packets moving in opposite directions, whose characteristics can be expressed in a universal way. Our treatment, which is hydrodynamic in nature, allows to describe the internal dynamics of these two packets in terms of the different velocities of their components. We confirm our analytical predictions through numerical simulations performed with the Gross-Pitaevskii equation and with the Calogero model (as an example of long range interactions and solvable with a parabolic confinement). Through the Calogero model we also discuss the effect of an external trapping on the protocol. The hydrodynamic approach shows that there is a difference between the bulk velocities of the propagating packets and the velocities of their peaks, accessible through different measurement procedures. -J. Phys. A: Math. Theor. 48 (2015) 28FT01; -New J. Phys. 18 (2016) 115003

¹H2020 Twinning project No. 692194, RBIT-WINNING; FP7 Marie Curie PIOF-PHY-276093; CUNY Award award 68193-00 46; European Project Matterwave

Fabio Franchini
Institut Ruer Bokovi

Date submitted: 06 Jan 2017

Electronic form version 1.4